Getting to the guts of SARS-CoV-2 infection





Immunofluorescent staining of SARS-CoV-2infected intestinal organoids. Nucleoprotein (NP) stains viral capsid. After 24 hours, single virus-infected cells are generally observed in organoids. These small infection clusters spread through the whole organoid after 60 hours. (Source: Lamers et al., 2020)

It has been well described how SARS-CoV-2 infects lung tissue and causes severe acute respiratory syndrome, but now there is evidence to emerge that the intestine may also be involved as a viral target. It is also known the SARS-CoV-2 receptor, ACE2 (angiotensin converting enzyme 2), is expressed on differentiated enterocytes. In the May 1 2020 edition of Science, Lamers et al show, using human small intestinal organoids (hSIOs), that enterocytes could be infected by SARS-CoV and SARS-CoV-2 and that significant titers of infectious viral particles could be detected in these cells. When the authors performed mRNA sequence analysis after viral infection of the hSIOs they identified a broad signature of cytokines and interferon stimulated genes (ISGs) attributed to type I and III interferon responses. The authors conclude that "intestinal epithelium supports SARS-CoV-2 replication, and hSIOs serve as an experimental model for coronavirus infection and biology".

Journal Article: Lamers et al., 2020. <u>SARS-CoV-2 productively</u> <u>infects human gut enterocytes</u>. Science

Article by Clive Gray

Also Read: Faecal shedding of SARS-CoV-2 by Kenneth Omollo